

REVIEW ARTICLE

Role of Immunopathology in Clinical Course of Malaria: A Review

Aye Aye Wynn^{1*}, Ohnmar Myint¹

¹ Department of Pathobiology and Medical Diagnostics, Faculty of Medicine and Health Sciences, Universiti Malaysia Sabah, Kota Kinabalu, Sabah, Malaysia

*Corresponding author's email: drwynnaa@ums.edu.my

Received: 27 March 2018

Accepted: 1 August 2018

Keywords: malaria, immunopathology, host, parasites

ABSTRACT

Malaria is a major health problem in various parts of the world especially affecting the tropical countries. It affects the vital organs causing severe complicated malaria. Clinical syndromes like severe cerebral anaemia, coagulation abnormalities, respiratory distress and severe anaemia can increase the mortality of malaria infected cases. Variation in individual susceptibility and severity and type of clinical presentations of malaria raises the need for study of both the parasite and host immune reactions as well as the contribution of inflammatory cytokines in malaria pathogenesis. This study explored the immunopathological basis and advances of severe malaria and their importance in pathogenesis of malaria and its complications. Previous and ongoing studies indicate that changes in endothelium during the sequestration of parasites in organs causes disruption of endothelial barrier function leading to serious effects of malaria. Parasite and host factors contribute to disturbance of cytokine regulation and escape of parasites from the immune system of the host. Immunopathological changes and dysregulation of cytokine production play central role in pathogenesis and disease severity in malaria.

INTRODUCTION

Malaria caused by the intracellular parasite *Plasmodium*, affects many people in the world especially those living in tropical countries. It affects the vital organs causing severe complicated malaria. Life-threatening complications such as cerebral malaria,

coagulation abnormalities, respiratory distress and severe anaemia can increase the mortality of malaria infected cases¹. Variation in individual susceptibility and severity as well as type of clinical presentations of malaria raises the need for study of both the parasite and host immunopathological mechanisms. Cytokines released by the host cells upon induction by parasite surface antigens play important role in tissue damage and red cell sequestration seen in severe malaria. Immunopathological basis of severe malaria and their importance in outcome prediction and success of management should be explored.

Endothelial Activation and Parasite Cytoadherence

Endothelial activation is a major pathogenetic mechanism in malaria pathogenesis. *Plasmodium falciparum* erythrocyte membrane protein-1 from parasites (PfEMP) is a product of diverse *var* gene²⁻⁵. PfEMP forms knobs on the parasitized red cell surface and binds ligands including CD36 and E-elastin on the endothelial cells which are then activated. Increased expression of adhesion molecules on endothelial surfaces occur and result in sequestration of red blood cells leading to ischaemia of the organ affected. Endothelial permeability is augmented by cytokine cascade³. The sequestration process results in firm adhesion of IEs to endothelial cells (ECs), monocyte recruitment, microcirculatory changes and induction of cytokine cascade causing local injury and dysfunction. Endothelial surface expression of intercellular adhesion molecule-1 (ICAM-1), endothelial protein C receptor (EPCR) and PECAM-1 are augmented in severe cases of *falciparum* malaria². These augment the inflammation around the minute vessels and lead to tissue and endothelial injury of pulmonary and brain microvasculature causing acute lung injury and disruption of blood brain barrier in cerebral malaria.

ICAM-1 and EPCR are receptors involved in cerebral malaria⁶. Studies showed that some haemoglobinopathies cause limited red cell invasion by the parasites. Haemoglobin S causes sickling of parasitized red cells rendering protection of malaria. Homozygous (HbSS) and heterozygous (HbAS) states have host microRNA (mRNA) profiles which after fusion with parasite mRNA render inhibition of parasite growth intracellularly. Host polymorphism that affects *P. falciparum* erythrocyte membrane protein-1 (PfEMP-1) may protect against malaria by impairing the parasite's ability to cytoadherence to microvessels⁷⁻⁹. Spleen is a major organ to remove malaria parasites from the circulation. Cytoadherence of malaria parasites is vital to the parasite survival to escape from splenic removal¹⁰. Virulence of the parasites differs according to the ability of cytoadherence through several parasite receptors such as plasmodium EMP1 (PfEMP1). PfEMP1 proteins mediate cytoadhesion of parasitized red cells lining cells of the organ microvasculature. pfEMP1 has dual binding specificity and these structures can be divided into group A (EPCR binders) and group B (CD 36 binders). Cerebral malaria is caused by dual binding of PfEMP to ICAM-1 and endothelial protein C receptor (EPCR) and this fact can be implicated in prevention of cerebral malaria in future¹¹⁻¹³.

Host Immunopathogenesis

In endemic areas of malaria, there is development of immunity by forming acquired antibodies against variant surface antigen (VSA) such as PfEMP, MSP3 and GLURP (RO)¹¹. Microparticles derived from platelets or parasitized RBC are seen in association with cerebral malaria in the sites of inflammation. Platelet-derived microparticles activate the capillary endothelium and regulate the pro-inflammatory cytokine production leading to increased vascular permeability^{14, 15}. Overproduction of TNF α , IL-1 β and chemokines induced by *plasmodium* glycoposphoinositol

(GPI) are responsible for disease mortality and deterioration^{6, 16}. Antibodies to GPI is closely correlated with parasitaemia and disease severity^{15 - 18}. Cytokine cascade is augmented by some chromosomal proteins called high mobility group box chromosomal protein 1 (HMGB1) which is secreted by activated mononuclear cells and passively through damaged cells. Levels of HMGB1 has been shown to parallel with disease severity and to induce permeability in endothelial cells, induce proinflammatory responses in macrophages through activation of TLR2, TLR4, or receptor for advanced glycation end products (RAGE)^{19, 20}. Elevated levels of HMGB1 can be used as a prognostic marker of disease severity in severe malaria^{20, 21}. IFN γ plays a crucial role in the clearance of intracellular pathogen by inducing the MHC molecules²². It also causes expression of gene encoding IDO (indoleamine 2, 3-dioxygenase), a rate limiting enzyme of tryptophan metabolism that can generate quinolinic acid (QA). Increased central level of QA is implicated in the causation of hyperexcitability, dementia and neurological dysfunctions seen in complicated malaria²³. CD40/CD40 ligand binding is important for binding of TNF activated platelets to the endothelial cells^{24, 25}. IL-1 increases the expression of ICAM1 and the production of cytokines (such as IL-6) by endothelial cells²⁴. Microparticles or moieties derived from blebbing of membranes of platelets and other cells during malaria infection. Platelet-derived microparticles can modulate the macrophage pro-inflammatory cytokine production and increase the endothelium permeability²⁶. Cell mediated immunity contributed by CD4+ T cells has a major role in immunity against malaria infection, both in pre-erythrocytic and erythrocytic stage^{27, 28}. They help to produce IFN γ and help B cells in control of malaria. People living in endemic areas of malaria possess IFN γ and IL-10 secreting CD4+ T cells²⁸.

Cytokines-Enhanced Haematological Abnormalities

Disseminated intravascular coagulation (DIC) is a life-threatening disorder occurring as a secondary to malaria. Expression of tissue factor (TF) is essential in initiation of blood coagulation. It occurs when the endothelial cells (EC) are exposed to pRBC. Initial stage of coagulation cascade after TF expression is escalated by amplification, propagation, and consolidation contributed by active role of sequestered pRBC and activated platelets at the sequestered sites²⁹. Severe anaemia in malaria can be caused by lysis of infected and uninfected RBCs, splenic sequestration of RBCs³⁰ dyserythropoiesis and bone marrow suppression³¹, erythrophagocytosis³² and chronic transmission of malaria in endemic regions. *P. falciparum*-derived haemozoin pigment (PfHz) and cytokines (TNF and IFN) promotes the host immune response and potentially causes suppression of the erythropoietic response³².

Role of Microglial Cells and Apoptosis in Malaria

Plasmodium apoptosis-linked pathogenicity factors (PALPF), PALPF-2, PALPF-5 can induce endothelial cell death in lining cells of microcapillaries in brain and lungs in severe malaria which are responsible for the development of acute respiratory distress and neurological abnormalities in severe malaria³³. CD8+ T cells act by direct cytotoxicity on endothelial cells by apoptosis or granzyme-induced lysis of cells. This can lead to disruption of blood-brain-barrier and development of cerebral malaria. Microglial cells are activated in human cerebral malaria and shown to produce matrix enzyme, metalloproteinase, and induce cytokines which can be applied in destruction of blood brain barrier and spread of infection to the central nervous system and neuron survival^{34, 35}.

Malaria Pigment: A Potential Prognostic Marker

Accumulation of haemozoin pigment (HZ) in the phagocytic cells of the immune system is used in the diagnosis and prognosis of malaria³⁶. *P. falciparum*-derived haemozoin pigment (PfHz) promotes the host immune response by activating NOD-like receptor of macrophages and potentially causes suppression of the erythropoietic response^{37, 38}. It can cause monocyte and macrophage dysfunction by impairing phagocytosis and the expression of MHC class II molecules and ICAM1, inhibiting dendritic cell (DC) maturation and proliferative responses by leucocytes³⁸.

Role of Nuclear Histones

Histones are acid-soluble proteins found in chromatin complexes released on rupture of parasites and host cells. Level of circulating histones in patients with falciparum malaria is correlated positively with disease severity³⁹. Histones can cause endothelial permeability and cytotoxicity by causing disruption of junctional proteins leading to cell death. Activation of toll like receptor (TLR2) and other receptors induces the release of IL-8 and other inflammatory mediators. Research is in progress to find out the potential uses of rhAPC that can cleave histones in hope to inhibit the cytokine induction and vascular permeability^{40, 41}.

Host Susceptibility

Susceptibility and severity of malaria infection is determined by a variety of host factors. Red blood cells carrying haemoglobin S (HbAS), HbAE, G6PD deficiency and alpha and beta thalassemia have reduced risk of developing severe anaemia by various protective effects such as reduced red cell invasion or impaired multiplication of parasites⁴². There are increased susceptibility and risk of severe malaria in individuals with polymorphism of adhesion molecules and cytokine such as

ICAM-1, PECAM1, TLR, CXCL10 and tumour necrosis factor (TNF)^{43–48}.

Host and Parasite Macrophage Inhibitory Factors (MIF)

Macrophage migration inhibitory factor (MIF) is a cytokine produced mainly by host macrophages. It regulates the expression of TNF α and inflammatory mediators such as nitric oxide and cyclooxygenase 2 (COX 2)⁴⁹. Plasmodium MIF (pMIF) is secreted when the parasites ruptured in schizont stage and they are exposed to immune cells. Levels of plasmodium MIF (pMIF) are positively correlated with parasitaemia, TNF α and IL-10. pMIF attenuates *Plasmodium* virulence by modulating functions of monocytes in host immune responses^{49–51}.

Vector-Parasite Association Affecting the Parasite Virulence

Studies have shown that vector mortality varies significantly among the different genotypes of parasites and environmental conditions⁵². Mosquitoes not only act as vectors but also modify the virulence of parasites. Transcriptomic studies showed after several blood passages, there is an expression of PIR gene in blood-stage parasites and increased virulence⁵³. Mosquito transmission modifies the diversity and magnitude of gene such as rifin and *var*⁵⁴ in malaria parasite which progress through each step of the lifecycle in both vector and host^{55, 56}.

CONCLUSION

Understanding of basic and advances in immunopathological processes that cause endothelial barrier dysfunction, sequestration of parasites, destructive effects of host and parasite factors and cytokine storm in malaria infection explains the need for defining clinical biomarkers of outcome. It also helps to identify possible new targets for management

in severe *falciparum* malaria such as trial of rhAPC to regulate the endothelial dysfunction and monoclonal anti-cytokine antibody or other drugs that block cytokine such as TNF to inhibit the activated macrophages.

REFERENCES

1. Marsh K, Kinyanjui S. (2006). Immune effector mechanisms in malaria. *Parasite Immunol* 28 (1 – 2): 51 – 60.
2. Jenkins NE, Chakravorty SJ, Urban BC, Kai OK, Marsh K, Craig AG. (2006). The effect of *Plasmodium falciparum* infection on expression of monocyte surface molecules. *Tropical Medicine and Hygiene* 100: 1007 – 1101.
3. Jenkins N, Wu Y, Chakravorty S, Kai O, Marsh K, Craig A. (2007). *Plasmodium falciparum* intercellular adhesion molecule-1- based cytoadherence-related signaling in human endothelial cells. *J Infect Dis* 15, 196 (2): 321 – 327.
4. Clark IA, Awburn MM, Harper CG, Liomba NG, Molyneux ME. (2003). Induction of HO-1 in tissue macrophages and monocytes in fatal *falciparum* malaria and sepsis. *Malar J*: 41.
5. Taylor TE, Fu WJ, Carr RA, Whitten RO, Mueller JS, Fos-iko NG, Lewallen S, Liomba NG, Molyneux ME. (2004). Differentiating the pathologies of cerebral malaria by post-mortem parasite counts. *Nat Med* 10: 143 – 145.
6. Ringwald P, Peyron F, Lepers JP, Rabarison P, Rakotomalala C, Razanamparany M, Rabodonirina M, Roux J, Le Bras J. (1993). Parasite virulence factors during *falciparum* malaria: Resetting, cytoadherence and modulation of cytoadherence by cytokines. *Infection and Immunity* 61 (12): 5198 – 5204.
7. Agarwal A, Guindo A, Cissoko Y, Taylor JG, Coulibaly D, Kone A, Kayentao K, Djimde A, Plowe CV, Doumbo O, Wellem TE, Diallo D. (2000). Hemoglobin C associated with protection from severe malaria in the Dogon of Mali, a West African population with a low prevalence of hemoglobin S. *Blood* 96: 2358 – 2363.
8. Williams TN. (2006). Human red blood cell polymorphisms and malaria. *Curr Opin Microbiol* 9: 388 – 394.
9. Guarin P, Primo L, Ferrandi C, Bussolino F, Tandon NN, Arese P, Ulliers D, Alessio M. (2001). Cytoadherence of *Plasmodium falciparum*-infected erythrocytes is mediated by a redox-dependent conformational fraction of CD36. *The Journal of Immunology* 167 (11): 6510 – 6517.
10. Chotivanich K, Udomsangpetch R, McGready R. (2002). Central role of the spleen in malaria parasite clearance. *The Journal of Infectious Diseases* 185: 1538 – 1541.
11. Avril M, Bernabeu M, Benjamin M, Brazier AJ, Smith JD. (2016). Interaction between endothelial protein C receptor and intercellular adhesion molecule 1 to mediate binding of *plasmodium falciparum*-infected erythrocytes to endothelial cells. *M Bio* 12, 7(4): e00615 – e00616.
12. Adams Y, Kuhnrae P, Higgins M K, Rowe JA. (2014). Resetting *Plasmodium falciparum*-infected erythrocytes bind to human brain microvascular endothelial cells in vitro, demonstrating a dual adhesion phenotype mediated by P *falciparum* erythrocyte membrane protein 1 domains. *Infect Immune* 82 (3): 949 – 959.
13. Rask TS, Hansen DA, Theander TG, Pedersen AG, Lavstse T. (2010). *Plasmodium falciparum* erythrocyte membrane protein 1 diversity in seven genomes – divide and conquer. *PLOS Computational Biology* 6 (9): 1 – 23. DOI: 10.1371/journal.pcbi.1000933.
14. Faille D, Combes V, Mitchell AJ, Fontaine A, Juhan-Vague I, Alessi M, Chimini G, Fusai T, Grau GE. (2009). Platelet microparticles: a new player in malaria parasite cytoadherence to human brain endothelium. *FASEB J* 23: 3449 -58.
15. Tamura T, Kimura K, Yuda M, Yui K. (2011). Prevention of experimental cerebral malaria by Flt3 ligand during infection with *Plasmodium berghei* ANKA. *Infect Immun* 79 (10): 3947 – 3956. DOI: 10.1128/IAI.01337-10.
16. Arrighi RB, Faye I. (2010) *Plasmodium falciparum* GPI toxin: A common foe for man and mosquito. *Acta Trop* 114 (3): 162 – 165.
17. Schofield L, Hewitt MC, Evans K, Siomos MA, Seeberger PH. (2002). Synthetic GPI as a candidate anti-toxic vaccine in a model of malaria. *Nature* 418: 785 – 789.

18. Naik RS, Branch OH, Amina S, Woods AS, Vijaykumar M, Perkins DJ, Nahlen BL, Lal AA, Cotter RJ, Costello CE, Ockenhouse CF, Davidson EA, Gowda DC. (2000). Glycosylphosphatidylinositol anchors of *Plasmodium falciparum*: Molecular characterization and naturally elicited antibody response that may provide immunity to malaria pathogenesis. *J Exp Med* 192: 1563 – 1576.
19. Wilson NO, Jain V, Roberts CE, Lucchi N, Joel PK, Singh MP, Nagpal AC, Dash AP, Udhayakumar V, Singh N, Stiles JK. (2011). CXCL4 and CXCL10 predict risk of fatal cerebral malaria. *Dis Markers* 230 (1): 39 – 49. DOI: 10.3233/DMA- pp 2011-0763.
20. Scaffidi P, Misteli T, Bianchi ME. (2002). Release of chromatin protein HMGB1 by necrotic cells triggers inflammation. *Nature* Jul 11, 418, (6894):191 – 195.
21. Higgins SJ, Xing K, Kim H, Kain DC, Wang F, Dhabangi A, Musoke C, Cserti-Gazdewich CM, Tracey KJ, Kain KC, Liles WC. (2012) Systemic release of high mobility group box 1 (HMGB1) protein is associated with severe and fatal *Plasmodium falciparum* malaria. *Malaria Journal* 12: 105.
22. Hunt NH, Ball HJ, Hansen AM, Khaw LT, Guo J, Bakmiwewa S, Mitchell AJ, Combes V, Grau GE. (2014). Cerebral malaria: Gamma-interferon redux. *Front Cell Infect Microbiol* 4: 11. DOI: 10.3389/fcimb.2014.00113 pp1-12.
23. Medana IM, Day NP, Salahifar-Sabet H, Stocker R, Smythe G, Bwanaisa L, Njobvu A, Kayira K, Turner GD, Taylor TE, Hunt NH. (2003). Metabolites of the kynurenine pathway of tryptophan metabolism in the cerebrospinal fluid of Malawian children with malaria. *JID* 188: 844 – 949.
24. Schofield L, Grau GE. (2005). Immunological processes in malaria pathogenesis. *Nature Reviews Immunology* 5: 722 – 735.
25. Piguet PF, Kan CD, Vesin C, Rochat A. (2001). Role of CD40-CD40L in mouse severe malaria. *Am J Pathol* 159 (2): 733 – 742.
26. Couper KN, Barnes T, Hafalla JCR, Combes V, Ryffel B, Secher T. (2010). Parasite-derived plasma microparticles contribute significantly to malaria infection-induced inflammation through potent macrophage stimulation. *PLOS Pathogens* 6 (1):1 – 13. DOI: 10.1371/journal.ppat.1000744
27. Gitau EN, James TJ, Karanja H, Stevenson L, Requena P, Kimani E, Olotu A, Kimani D, Marsh K, Bull P, Urban BC. (2014). CD4+ T cell responses to *plasmodium falciparum* erythrocyte membrane protein 1 in children with mild malaria. *J Immunol* 192 (4): 1753 – 1761. Available at [http:// www.jimmunol.org/content/early/2014/01/22/jimmunol.1200547](http://www.jimmunol.org/content/early/2014/01/22/jimmunol.1200547).
28. Perez-Malich D, Langhorne J. (2015). CD4-T cell subsets in malaria: TH1/TH2 revisited. *Front Immunol* 5: 671.
29. Francischetti IMB, Seydel KB, Monteiro RQ. (2008). Blood coagulation, inflammation and malaria. *Microcirculation* Feb 15 (2): 81 – 107. DOI: 10.1080/10739680701451516
30. Imamura T, Sugiyama T, Cuevas LE, Makunde R, Nakamura S. (2002). Expression of tissue factor the clotting initiator, on macrophages in *plasmodium falciparum* infected placentas. *J Infect Dis* 186 (3): 436 – 440.
31. Helleberg M, Goka BQ, Akanmori BD, Obeng-Adjei G, Rodrigues O, Kurtzthals. (2005). Bone marrow suppression and severe anaemia associated with persistent *Plasmodium falciparum* infection in African children with microscopically undetectable parasitaemia. *Malar J* 4 (56): 1 – 7.
32. Arese P, Turrini F, Ginsburg H. (1991). Erythrophagocytosis in malaria: Host defence or menace to the macrophage? *Parasitology Today* 7 (1): 123 – 128.
33. Nadine N, Dilimabaka N, Taoufiq Z, Zougbede S, Bonnefoy SM, Lorthiosis A, Couraud PO, Rebollo A, Snounou G, Mazier D, Sabater AM. (2014). *P. falciparum* isolate-specific distinct patterns of induced apoptosis in pulmonary and brain endothelial Cells. *PLoS ONE* 9 (3): 2014, e90692. DOI:10.1371/journal.pone.0090692
34. Mariani MM, Kielian T. (2009). Microglia in infectious disease of the central nervous system. *J Neuroimmune Pharmacol* 4 (4): 448 – 446.
35. Schcluesener H, Kreamsner P, Meyermann R. (1998). Widespread expression of MRP-8 and MRP14 in human cerebral malaria by microglial cells. *Acta Neuropathol* 96: 575 – 580.
36. Olivier M, Van Den Ham K, Shio MT, Kassa FA, Fougeray S. (2014). Malarial pigment hemozoin and the innate inflammatory response. *Front Immunol* 5: 25. DOI: 10.3389/fimmu.2014.00025.

37. Perkins DJ, Were T, Davenport GC, Kempaiah P, Hittner JB, Ong'echa JM. (2011). Severe malarial anemia: Innate immunity and pathogenesis. *Int J Biol Sci* 2011, 7 (9): 1427 – 1442. DOI:10.7150/ijbs.7.1427.
38. Dong Liu, Rhebergen AM, Stephanie C, Eisenbarth SC. (2013). Licensing adaptive immunity by NOD-like receptors. *Front Immunol* 4: 486. DOI: 10.3389/fimmu.2013.00486
39. Mark R, Gillrie MR, Lee KD, Gowda C, Davis SP. (2012). *Plasmodium falciparum* histones induce endothelial proinflammatory response and barrier dysfunction. *Immunopathology and Infectious Diseases. Am J Pathol* 180: 1028 – 1039. DOI: 10.1016/j.ajpath.2011.11.037
40. Xu J, Zhang X, Pelayo R, Monestier M, Ammollo CT, Semeraro F, Taylor FB, Esmon NL, Lupu F, Esmon CT. (2009). Extracellular histones are major mediators of death in sepsis. *Nat Med* 15: 1318 – 1322.
41. Monal Sharma, Chhaya Dhiman, Poonam Dangi, Shailja Singh. (2014). Designing synthetic drugs against *Plasmodium falciparum*: A computational study of histone-lysine N-methyltransferase (PfHKMT). *Syst Synth Biol* 8: 155 – 160. DOI: 10.1007/s11693-014-9144-8.
42. Min OG, Gros P. (2005). Erythrocyte variants and the nature of their malaria protective effect. *Cellular Microbiology* (6): 753 – 763.
43. Sinha S, Qidwai T, Kanchan K, Anand P, Jha GN, Pati SS, Mohanty S, Mishra SK, Tyagi PK, Sharma SK. (2008). Variations in host genes encoding adhesion molecules and susceptibility to falciparum malaria in India. *Malaria J* 7 (250):1 – 9.
44. de Mendonça VRR, Goncalves MS, Barral-Netto M. (2012). The host genetic diversity in malaria infection. *Journal of Tropical Medicine* 1 – 17. Available at <http://dx.doi.org/10.1155/2012/940616>
45. Apinjoh TO, Anchang-Kimbi JK, Njua-Yafi C, Mugri RN, Ngwai AN, Rockett KA, Mbunwe E, Besingi RN, Clark TG, Kwiatkowski DP, Achidi EA. (2013). Association of cytokine and toll-like receptor gene polymorphisms with severe malaria in three regions of Cameroon. *PLoS ONE* 8 (11): e81071. Available at <https://doi.org/10.1371/journal.pone.0081071>
46. Gichohi-Wainaina WN, Melse-Boonstra A, Feskens EJ, Demir AY, Veenemans J, Verhoef H. (2015). Tumour necrosis factor allele variants and their association with the occurrence and severity of malaria in African children: A longitudinal study. *Malaria Journal* 14 (249): 1 – 11. DOI: 10.1186/s12936-015-0767-3
47. Wilson N, Driss A, Solomon W, Dickinson-Copeland C, Salifu H, Jain V, Singh N, Stiles J (2013). CXCL10 Gene Promoter Polymorphism -1447A>G Correlates with Plasma CXCL10 Levels and is associated with male Susceptibility to cerebral malaria. *PLoS ONE* 8 (12): e81329. DOI: 10.1371/journal.pone.0081329
48. Mockenhaupt FP, Cramer JP, Hamann L, Stegemann MS, Eckert J, Oh NR, Otchwemah RN, Dietz E, Ehrhardt S, Schröder NWJ, Bienzle U, Ralf R, Schumann RR. (2006). Toll-like receptor (TLR) polymorphisms in African children: Common TLR-4 variants predispose to severe malaria. *Proceedings of the National Academy of Sciences of the United States of America* 103 (1): 177 – 182.
49. Rosado JD, Rodriguez-Sosa M. (2011). Macrophage migration factor (MIF): A key player in protozoan infections. *Int J Biol Sci* 7 (9): 1239 – 1256. DOI:10.7150/ijbs.7.1239
50. Han C, Lin Y, Shan G, Zhang Z, Sun X, Wang Z, Wei C, Deng Y, Zhang L, Bu L, Shao D, Wang H. (2010). Plasma concentration of malaria parasite-derived macrophage migration inhibitory factor in uncomplicated malaria patients correlates with parasitemia and disease severity. *Clin Vaccine Immunol* 17 (10): 1524 – 1532.
51. Bozza MT, Martins YC, Carneiro LAM, Paiva CN. (2012). Macrophage migration inhibitory factor in protozoan infections. *Journal of Parasitology Research*, Article ID 413052, 12.
52. Ferguson HM, Read AF. (2002). Genetic and environmental determinants of malaria parasite virulence in mosquitoes. *Proc R Soc Lond B* 269: 1217 – 1224.
53. Lee HJ, Georgiadou A, Otto TD, Levin M, Coin LJ, Conway DJ, Cunningham AJ. (2018). Transcriptomic studies of malaria: A paradigm for investigation of systemic host-pathogen interaction. *Microbiol Mol Biol Rev.* 82 (2) e00071-17: 1 – 37.

54. Mackinnon MJ. (2014). The role of immunity in mosquito-induced attenuation of malaria virulence. *Malar J* 13: 25. DOI: 10.1186/1475-2875-13-25. pmid:24443873
55. Florens L, Washburn MP, Raine JD, Anthony RM, Grainger M, Haynes JD, et al. (2002). A proteomic view of the *Plasmodium falciparum* life cycle. *Nature* 419 (6906): 520 – 526. pmid:12368866. DOI: 10.1038/nature01107
56. Spence PJ, Brugat T, Langhorne J. (2015). Mosquitoes reset malaria parasites. *PLoS Pathog* 11 (7): 1 – 5, e1004987. DOI:10.1371/journal.ppat.1004987